

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

Form Approved
OAR No. 0704 01AA

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION

Unclassified

2a SECURITY CLASSIFICATION AUTHORITY

2b DECLASSIFICATION/DOWNGRADING SCHEDULE

4 PERFORMING ORGANIZATION REPORT NUMBER(S)

6a NAME OF PERFORMING ORGANIZATION

Purdue University

6c ADDRESS (City, State, and ZIP Code)

West Lafayette, IN 47907

8a NAME OF FUNDING/SPONSORING
ORGANIZATION

Office of Naval Research

8c ADDRESS (City, State, and ZIP Code)

800 North Quincy Street
Arlington, VA 22217-50006b OFFICE SYMBOL
(If applicable)8b OFFICE SYMBOL
(If applicable)

1b RESTRICTIVE MARKINGS

3 DISTRIBUTION/AVAILABILITY OF REPORT

Distribution Lists Enclosed

5 MONITORING ORGANIZATION REPORT NUMBER(S)

7a NAME OF MONITORING ORGANIZATION

Office of Naval Research

7b ADDRESS (City, State, and ZIP Code)

Department of the Navy
Arlington, VA 22217

9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-89-J-1128

10 SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NOPROJECT
NOTASK
NOWORK UNIT
ACCESSION NO

11 TITLE (Include Security Classification)

Chiral Oxazaborolidine as Catalysts for the
Enantioselective Addition of Diethylzinc to Aldehydes

12 PERSONAL AUTHOR(S)

N. N. Joshi, M. Srebnik and H. C. Brown

13a TYPE OF REPORT

Technical

13b TIME COVERED
FROM TO

14 DATE OF REPORT (Year, Month, Day)

July 20, 1989

15 PAGE COUNT

10

16 SUPPLEMENTARY NOTATION

17 COSATI CODES

| FIELD | GROUP | SUB GROUP |
|-------|-------|-----------|
| | | |
| | | |
| | | |

18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Chiral Oxazaborolidines, Ephedrine, Diethylzinc,
Enantioselective Addition, Chiral *sec*-alcohols

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

DTIC
ELECTE
S JUL 26 1989 **D**
D

see attached report

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

20 DISTRIBUTION/AVAILABILITY OF ABSTRACT

☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS

22a NAME OF RESPONSIBLE INDIVIDUAL

Professor Herbert C. Brown

21 ABSTRACT SECURITY CLASSIFICATION

Unclassified

22b TELEPHONE (Include Area Code)

(317) 494-5316

22c OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

89 . 7 26 052

AD-A210 543

OFFICE OF NAVAL RESEARCH

Contract N00014-89-J-1128

R & T code 4135011---04

Technical Report No. 3

Chiral Oxazaborolidines as Catalysts for the Enantioselective
Addition of Diethylzinc to Aldehydes

by

N. N. Joshi, M. Srebnik, and H. C. Brown

Prepared for Publication

in

Tetrahedron Letters

Purdue University
H. C. Brown and R. B. Wetherill Laboratories of Chemistry
West Lafayette, Indiana 47907

July 20, 1989

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

* This document has been approved for public release
and sale; its distribution is unlimited.

PURDUE UNIVERSITY



July 21, 1989

HERBERT C. BROWN

Professor Harry Wasserman, Editor
Tetrahedron Letters
Department of Chemistry
Yale University
P. O. Box 6666
New Haven, Connecticut 06511

Dear Harry:

Enclosed are the copyright, two original camera-ready copies, and two graphical abstracts of a manuscript entitled, *Chiral Oxazaborolidines as Catalysts for the Enantioselective Addition of Diethylzinc to Aldehydes*, by N. N. Joshi, M. Srebnik and H. C. Brown, hereby submitted for publication in Tetrahedron Letters.

Cordially,

A handwritten signature in cursive script, appearing to read "Herb", followed by a horizontal line.

Herbert C. Brown

HCB:ms
encl.

cc: N. N. Joshi
M. Srebnik



H. Guard
I. Klundt
C. F. Lane

Chiral Oxazaborolidines as Catalysts for the Enantioselective Addition of Diethylzinc to Aldehydes.

N. N. Joshi, M. Srebnik and Herbert C. Brown

H. C. Brown and R. B. Wetherill Laboratories of Chemistry, Purdue University, West Lafayette, Indiana 47907 USA.

The oxazaborolidine, **1**, conveniently derived from (-)-ephedrine and $\text{BH}_3\cdot\text{SMe}_2$ provides an excellent catalyst for the enantioselective addition of diethylzinc to aromatic aldehydes. This is the first example of a chiral boron intermediate for this reaction.



| | |
|-----------------|--------|
| Accession | |
| NTIS | CRAM ✓ |
| DTIC | TAB 11 |
| Univ. of Calif. | 11 |
| Joshi | |
| By | |
| Date | |
| Title | |
| Author | |
| Subject | |
| A-1 | |


Chiral Oxazaborolidines as Catalysts for the Enantioselective Addition of Diethylzinc to Aldehydes.

N. N. Joshi, M. Srebnik* and Herbert C. Brown*

H. C. Brown and R. B. Wetherill Laboratories of Chemistry, Purdue University, West Lafayette, Indiana 47907 USA

Summary. (4*S*, 5*R*)-4-methyl-5-phenyl-1,3,2-oxazaborolidine, **1**, catalyzes the addition of diethylzinc to aldehydes to furnish secondary alcohols in up to 96% ee. This is the first example of a chiral boron intermediate for this reaction.

While the chemistry involving stoichiometric reactions of organoboranes has been amply investigated and shown to be rich and diversified, catalytic processes especially asymmetric, have been relatively neglected, and only recently have a number of enantioselective reactions catalyzed by chiral organoboranes appeared in the literature. However, since many reactions are catalyzed by Lewis acids and since chiral organoboranes are readily accessible chiral Lewis acids, we have undertaken a program to develop new catalysts based on boron and to apply them to suitable reactions. The addition of diethylzinc, DEZ, to aldehydes is one such reaction. DEZ reacts with aldehydes very sluggishly at 25°C in non-coordinating solvents³. In 1984 Oguni and Omi reported that small amounts of amino alcohols greatly accelerated the reaction and induced optical activity, albeit modest, when optically active amino alcohols were involved (eq 1).



Subsequent workers increased the enantiomeric excess (ee) of the product alcohols but only at the expense of designing more and more complex amino alcohols⁵. Examination of the postulated mechanism of this reaction indicated to us that boron might well serve as the central metal in the template and by virtue of generally forming shorter B-O and B-N bonds than do other metals, possibly provide a superior catalytic system. In this communication we describe our initial efforts into asymmetric catalysis with chiral organoboranes in the particular case of the addition of DEZ to aldehydes. As structures for possible catalysts, we chose the oxazaborolidines⁶. As the amino alcohol moiety we selected ephedrine, both enantiomers of which are commercially available. The borolidines were prepared by heating the amino alcohol with either $\text{BH}_3 \cdot \text{SMe}_2$ or the appropriate boronate ester, $\text{R}(\text{BOR}')_2$ until two equivalents of either H_2 or $\text{R}'\text{OH}$ had evolved (eq 2).





Table 1. Reaction of Diethylzinc with Benzaldehyde Catalyzed by Chiral Oxazaborolidines^a.



| Entry | Config. of Catalyst | R | Time, h | % ee | Config. of Product |
|-------|---------------------|----|---------|------|--------------------|
| 1 | 4S, 5R | Ph | 60 | 47 | R |
| 2 | 4S, 5R | Me | 24 | 78 | R |
| 3 | 4S, 5R | H | 12 | 86 | R |
| 4 | 4R, 5S | H | 12 | 85 | S |
| 5 | 4S, 5S | H | 8 | 2 | R |
| 6 | b | b | 12 | 58 | R |
| 7 | c | c | 4 | 70 | R |
| 8 | d | d | 4 | 71 | R |

^a Reactions were run at 0°C in toluene, 3 M in DEZ, in the presence of 5 mol % catalyst. ^b Catalyst prepared by mixing AlMe₃ and (-)-ephedrine. ^c Catalyst prepared from *n*-BuLi and (-)-ephedrine. ^d Catalyst prepared by mixing DEZ and (-)-ephedrine.

The results for the addition of DEZ to benzaldehyde for several catalysts are summarized in Table 1. For the sake of comparison, we also prepared catalysts based on Al, Li and Zn by reacting ephedrine with AlMe₃, *n*-BuLi and DEZ respectively. (Presumably, these catalysts contain Al, Li, or Zn atoms in place of boron. However, since the exact structure of these catalysts in solution are not known their representation at this point is merely speculative.) A comparison of the results obtained with these metallated oxaza-ring systems indicates that in the present case the results with Al (58% ee), Li (70% ee), and Zn (71% ee) are clearly inferior to that obtained with boron (86% ee). As regards the present series of oxazaborolidines the following generalizations apply: 1. Both the rate and the ee are optimized when boron is unhindered (Table 1.). 2. The *cis* configuration of C4/C5 is preferred to *trans*. Based on these results we selected **1**, or its enantiomer, for further study. **1** exists essentially as a monomer in solution (1 M in toluene) as determined by ¹¹B-NMR. To ascertain the stability of **1**, (and the other oxazaborolidines in the presence of DEZ), we added an equivalent of DEZ to **1**. No discernible change in the ¹¹B-NMR spectrum was observed for 24 h at 25°C, indicative that the borolidine ring had not been cleaved or that exchange between BH and DEZ had not occurred⁷.

Aromatic aldehydes generally reacted with DEZ much faster than did aliphatic aldehydes. The optical inductions were good to excellent for aromatic aldehydes while the aliphatic aldehyde tested gave a product of only moderate ee⁸. Typical results are listed in Table 2.

Table 2. Reaction of DEZ and Various Aldehydes in the Presence of 1^a.

| Entry | Aldehyde | Time, h | $[\alpha]_D, c=5, \text{CHCl}_3$ | % ee | Abs. Config. |
|-------|-----------------------|---------|----------------------------------|-----------------|----------------|
| 1 | Benzaldehyde | 30 | +39.1 | 95 ^b | R ^d |
| 2 | 1-Methoxybenzaldehyde | 48 | +44.5 | 91 ^b | R ^e |
| 3 | 4-Methoxybenzaldehyde | 48 | +32.2 | 93 ^b | R ^f |
| 4 | 4-Chlorobenzaldehyde | 48 | +27.7 | 96 ^b | R ^f |
| 5 | 2-Furaldehyde | 64 | +16.1 | 66 ^c | R ^g |
| 6 | Heptanal | 88 | -4.3 | 52 ^c | R ^f |

^a All reactions were run at -25°C in toluene, 3 N in DEZ and 5 mol% catalyst. Chemical yields in all cases were $\geq 85\%$. ^b Determined as the (-)-menthyl chloroformate ester on a Supelcowax capillary column, 15 m. ^c Determined as the (+)-MTPA ester on a SPB-5 capillary column, 30 m. ^d Ref. 4. ^e Ref. 5a. ^f Ref. 5f. ^g Tentatively assigned by analogy to the other compounds.

As regards the mechanism of this reaction induced by oxazaborolidines, we believe that an intermediate ternary complex is formed, consisting of the aldehyde complexed to boron and DEZ coordinated to either nitrogen or oxygen of the borolidine ring⁹. In addition since no precatalyst-catalyst equilibrium is operating in the present case as it is with the zinc derived catalyst¹⁰, a linear relationship between the optical purity of the catalyst and the ee of the product should exist¹¹. Indeed, we have found this to be the case, again indicative that a single monomeric oxazaborolidine species is involved in the catalysis.

In conclusion, this is the first example of the use of a chiral boron intermediate as a catalyst for the addition of an organometallic reagent to aldehydes. We are currently probing the precise mechanism of this reaction, an understanding of which should enable us to devise superior catalysts capable of handling both aromatic and aldehydes. Other asymmetric transformations amenable to catalysis by organoboranes are also being investigated and will be the subject of future publications.

Acknowledgement: This work was supported in part by The Office Of Naval Research to whom we express our appreciation.

References and Notes

1. Pelter, A.; Smith, K.; Brown, H. C. *Boron Reagents*; Academic Press: London, 1988.
2. **Diels-Alder reaction:** a. Furuta, K.; Shimizu, S.; Miwa, Y.; Yamamoto, H. *J. Org. Chem.* **1989**, *54*, 1483. b. Furuta, K.; Miwa, Y.; Iwanaga, K.; Yamamoto, H. *J. Am. Chem. Soc.* **1988**, *110*, 6254. c. Bir, G.; Kaufmann, D. *Tetrahedron Lett.* **1987**, *28*, 777. **Silylcyanation** d. Reetz, M. T.; Kunisch, F.; Heitmann, P. *Tetrahedron Lett.* **1986**, *27*, 4721. **Reduction:** e. Itsuno, S.; Sakurai, Y.; Ito, K.; Hirao, A.; Nakahama, S. *Bull. Chem. Soc. Jpn.* **1987**, *60*, 395. f. Corey, E. J.; Bakshi, R. K.; Shibata, S. *J. Am. Chem. Soc.* **1987**, *109*, 5551.
3. Miginiac, L. In *The Chemistry of the Metal-Carbon Bond*; Hartley, F. R., Ed.; Wiley: Chichester, 1985; Vol. 1.
4. Oguni, N.; Omi, T. *Tetrahedron Lett.* **1984**, *25*, 2823.

5. a. Soai, K.; Niwa, S.; Watanabe, M. *J. Chem. Soc. Perkin Trans I* **1989**, 109. b. Yoshioka, M.; Kawakita, T.; Ohno, M. *Tetrahedron Lett.* **1989**, *30*, 1657. c. Oppolzer, W.; Radinov, R. N. *Tetrahedron Lett.* **1988**, *29*, 5645. d. Itsuno, S.; Frechet, J. M. J. *J. Org. Chem.* **1987**, *52*, 4140. e. Corey, E. J.; Hannon, F. J. *Tetrahedron Lett.* **1987**, *28*, 5237. f. Soai, K.; Ookawa, A.; Kaba, T.; Ogawa, K. *J. Am. Chem. Soc.* **1987**, *109*, 7111. g. Kitamura, M.; Suga, S.; Kawai, K.; Noyori, R. *J. Am. Chem. Soc.* **1986**, *108*, 6071. h. Smaardijk, Ab. A.; Wynberg, H. *J. Org. Chem.* **1987**, *52*, 135. i. Muchow, G.; Vannoorenberghe, Y.; Buono, G. *Tetrahedron Lett.* **1987**, *28*, 6163.

6. Barton, L. In *Gmelin: Handbuch der Anorganischen Chemie*; Springer Verlag: Berlin, 1977; Vol. 48, Supplement 16, p 101.

7. However, we have discovered that various boron esters do exchange with DEZ, the rate of exchange being proportional to the steric bulk of the alkoxy unit. Thus, B-methoxy-9-BBN completely exchanges with DEZ within 5 min at 25° C, each of the reactants being 1 M in toluene, while B-isopropoxy-9-BBN requires 3-4 h under the same conditions. B-t-butoxy-9-BBN does not exchange for at least 24 h. For a general treatise on the chemistry of dialkylzincs see: Boersma, J. In *Comprehensive Organometallic Chemistry*; Wilkinson, G. Ed.; Pergamon Press: Oxford, 1982; Vol. 2, p 823.

8. **Preparation of oxazaborolidine 1.** A solution of (1*R*, 2*S*)-(-)-ephedrine (8.25 g, 50 mmol) in anhydrous THF (50 mL) was treated with boron-methyl sulfide, BMS, (50 mmol, 5 mL of 10 M solution). The reaction mixture was stirred at 25° C for 1 h by which time one equiv of hydrogen had evolved. The volatiles were pumped off to furnish a white solid, ¹¹B-NMR (δ ~ 8 ppm). The solid was gradually heated to 100°C and maintained at that temperature until the second equivalent of hydrogen had evolved. The product was distilled under reduced pressure to obtain pure **1**, as a colorless liquid. Yield: 7.6 g (86 %); bp 110-112° C, 20 mm Hg; ¹¹B-NMR δ 25 ppm, d (J=160 Hz); IR_{BH} 2562 cm⁻¹. **The preparation of (R)-(+)-1-phenyl-1-propanol is typical:** A solution of diethylzinc (12 mmol, 4 mL) in toluene was cooled to -25° C using a cryobath. The catalyst **1**, (0.5 mmol, 1 mL of a 1 M solution) in toluene and benzaldehyde (10 mmol, 1.06 g) were then successively introduced into the reaction flask. A yellow color developed which gradually disappeared in the course of the reaction. A positive pressure of Ar was maintained inside the flask and the reaction was periodically monitored by GC (every 8 h), by which procedure the reaction was found to tail-off towards the end. (This could be prevented by using a larger excess of DEZ, ~ 50 %). At this stage the reaction mixture was diluted with ether (100 mL), transferred to a separatory funnel and cautiously washed with 2 N HCl followed by saturated brine. The organic phase was dried over anhydrous K₂CO₃ and concentrated under reduced pressure. The residue was purified by flash chromatography followed by short-path distillation to furnish (R)-(+)-1-phenyl-1-propanol. Yield: 1.2 g. (88 %); bp 105-107 °C, 20 mm Hg. The analysis for the enantiomeric excess was performed via the (-)-MCF ester on a 15 m Supelcowax capillary column: temp., 180°C, head pressure, 6 psi; R_t (for the major isomer) 10.8 min, 95% ee.

9. The steric environment around nitrogen and oxygen presumably determines the extent of coordination of DEZ with one or the other and in turn influences the magnitude of the optical induction. Unpublished observations.

10. Kitamura, M.; Okada, S.; Suga, S.; Noyori, R. *J. Am. Chem. Soc.* **1989**, *111*, 4028.

11. Oguni, N. Matsuda, Y.; Kaneko, T. *J. Am. Chem. Soc.* **1988**, *110*, 7877.

TECHNICAL REPORT DISTRIBUTION LIST, GENERAL

| | <u>No. Copies</u> | | <u>No. Copies</u> |
|--|------------------------------|--|-----------------------|
| Office of Naval Research Chemistry Division, Code 1113 800 North Quincy Street Arlington, VA 22217-5000 | 3 | Dr. Ronald L. Atkins Chemistry Division (Code 385) Naval Weapons Center China Lake, CA 93555-6001 | 1 |
| Commanding Officer Naval Weapons Support Center Attn: Dr. Bernard E. Doua Crane, IN 47522-5050 | 1 | Chief of Naval Research Special Assistant for Marine Corps Matters Code 00MC 800 North Quincy Street Arlington, VA 22217-5000 | 1 |
| Dr. Richard W. Drisko Naval Civil Engineering Laboratory Code L52 Port Hueneme, California 93043 | 1 | Dr. Bernadette Eichinger Naval Ship Systems Engineering Station Code 053 Philadelphia Naval Base Philadelphia, PA 19112 | 1 |
| Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314 | 2 <u>high quality</u> | Dr. Sachio Yamamoto Naval Ocean Systems Center Code 52 San Diego, CA 92152-5000 | 1 |
| David Taylor Research Center Dr. Eugene C. Fischer Annapolis, MD 21402-5067 | 1 | David Taylor Research Center Dr. Harold H. Singerman Annapolis, MD 21402-5067 ATTN: Code 283 | 1 |
| Dr. James S. Murday Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000 | 1 | | |